

# Tsama Mix

## DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- To be used only where there is a recognized need.
- Do not exceed the appropriate application rates.
- Avoid application under climate extremes.
- Wear suitable protective clothing.
- Clean all equipment before and after use.
- To be used only when recommended by a suitably qualified person.

### Mixing instructions:

Always follow the label recommendations. Where water volume differ from typical rates as specified on the label, adjust application rates accordingly.

Shake container thoroughly before use. Pour through the top filter into ¼ filled tank. Keep agitation going. Top up with water and continue agitating whilst spraying.

### Compatibility:

Read all labels carefully. This product may be compatible with many agrochemicals. Check with your supplier before tank mixing or contact Tsama Agricultural Agencies.

### Storage:

Keep in a cool dry chemical store, out of reach of children and livestock. Protect from frost and heat extremes.

### Conditions of Sale:

The registration holder warrants that the product within the unopened container conforms technically to the description on this label. The directions on this label must be followed carefully. As the use of this product is beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of use or results obtained. Use of the product implies purchaser accepts these conditions of sale.

### Recommendations:

Product can be applied either as a tractor spray or through the irrigation system. Water rate for tractor spray is recommended at 200 - 500 l/ha and through the irrigation at 100 000 l/ha.

Where TRV's differ from those on the label, adjust application rates per 100 l water accordingly so that the product quantity per ha applied remains the same. However, do not concentrate product up by more than 0.4%.

CROP	DOSAGE RATE	TIMING
Field crops & vegetables	0.5 - 3 kg/ha per application but not exceeding 6 kg/ha per season	Apply as soon as there is enough leaf material to intercept spray. Repeat at 10 - 14 day intervals if necessary.
Maize	1 - 3 kg/ha per application but not exceeding 6 kg/ha per season	Apply at 4 - 6 leaf stage and repeat at 6 - 8 leaf stage if necessary.
Potatoes	1 - 2 kg/ha per application to apply max 6 kg/ha per season	Apply 1 week after 100 % emergence and repeat at tuber initiation and 2 weeks later if necessary.
Lucerne & pastures	1 - 3 kg/ha but not exceeding 6 kg/ha per season	Apply after each cut or grazing when there is enough leaf material to absorb spray (typically a plant height of 10 - 15 cm). Ensure 14 days between application and grazing.
Onions	1 - 2 kg/ha per application but not exceeding 6 kg/ha per season	Apply when plant is 10 - 15cm high and repeat at bulb formation if necessary.
Macadamias & nut crops	Apply typically 100 g up to 200 g/100 l water based on TRV of 1 000 l/ha to apply 1 - 2 kg/ha per application but not exceeding a total of 6 kg/ha per season	Apply post harvest, before flowering. Repeat at 80 % petal fall if necessary.
Citrus	Apply typically 20g/100 l up to a max of 40 g/100 l water based on TRV of 6 000 l/ha to apply typically 1 kg/ha per application and a maximum of 2 kg/ha per application but not exceeding 6 kg/ha per season	Apply post harvest, before flowering and repeat at 80% petal fall if necessary.
Subtropical fruit	Apply typically 100 g/100 l to max 200 g/100 l water based on TRV of 1 000l/ha to apply typically 1 kg to a maximum of 2 kg/ha per application but not exceeding 6 kg/ha per season	Apply post harvest, before flowering and repeat at 80% petal fall if necessary.
Stone fruit	Apply typically 100 g/100 l to max 200 g/100 l water based on TRV of 1 000l/ha to apply typically 1 kg to a maximum of 2 kg/ha per application but not exceeding 6 kg/ha per season	Apply post harvest and at pink bud or at 80% petal fall if necessary.
Tomatoes & other horticultural crops	Apply 0.5 - 3 kg/ha per application, not exceeding 6 kg/ha per season	Apply from 7 days post transplant and repeat at 7 - 10 day intervals if necessary.
Vines	1 kg/ha at a maximum spray concentration of 0.1 %. Do not exceed 4 kg/ha per season.	Apply between bud break and first flowers visible or post harvest. Repeat applications may be applied at 10 - 14 day intervals if necessary.
Wheat	1 - 3 kg/ha per application. Do not exceed a total of 6 kg/ha per season.	Apply at 4 - 6 leaf stage and repeat at 10 - 14 day intervals if necessary with the last application before flag leaf visible.